

- ORNAMENTALS (continued)**
- Gypsophila (1 to 7)
  - Hawthorn (1 to 7)
  - Holly (1 to 7)
  - Hoya (1 to 7)
  - Hydrangea (1 to 7)
  - Impatiens\* (1 to 7)
  - Iris (1 to 8)
  - Juniper (1 to 7)
  - Kalanchoe (1 to 7)
  - Lilies (1 to 7)
  - Lipstick vine (1 to 7)
  - (Aeschynanthus)
  - Marigold (1 to 7)
  - Monarda (Bee Balm) (1 to 7)
  - Pachysandra (1 to 7)
  - Palm (1 to 7)
  - Pansy (1 to 7)
  - Peach (ornamental) (1 to 7)
  - Peperomia (1 to 7)
  - Periwinkle (1 to 7)
  - Philodendron (1 to 7)
  - Phlox (1 to 7)
  - Pilea (1 to 7)
  - Pine (1 to 7)
  - Pitosporum (1 to 7)
  - Plum (ornamental) (1 to 7, 14)
  - Poinsettia (1 to 7)
  - Poppy (1 to 7)
  - Monarda (Bee Balm) (1 to 7)
  - Pachysandra (1 to 7)
  - Privet (1 to 7)
  - Protea (1 to 7)
  - Pyracantha (1 to 7)
  - Rhododendron (1 to 7, 16)
  - Rose Tree of China (1 to 7)
  - Rose (1 to 7, 15)
  - Salvia (1 to 7)
  - Schefflera (1 to 7)
  - Snappdragon (1 to 7)
  - Statice (1 to 7)
  - Tree Ivy (1 to 7)
  - Tulip (1 to 7, 9)
  - Viburnum (1 to 7)
  - Violet (1 to 7)
  - Zinnia (1 to 7, 10)

\*NOTE: Do not use 26 GT Fungicide as a soil drench on Impatiens or Pothos. Do not use 26 GT Fungicide on Spathiphyllum.

**HOW TO USE 26 GT FUNGICIDE AS A FOLIAR SPRAY**

When conditions become favorable for disease development, apply 26 GT Fungicide as a foliar spray to insure thorough coverage of the plant foliage at the following rates and intervals:

DISEASE	QUARTS PRODUCT/ACRE	INTERVAL OF APPLICATION	USE DIRECTIONS
Aerial Web Blight Alternaria Leaf Blight Alternaria Leaf Blight Alternaria Leaf Spot	1.0 to 2.5	7 to 14 Days	Spray plants to insure thorough coverage.
Botrytis Blight Fusarium Leaf Spot Helminthosporium Leaf Spot Ink Spot Ray Blight Tulip Fire Daffodil Leaf Scorch Blossom Blight			

Do not apply more than 2.5 quarts product/acre per application. Do not make more than 4 applications per crop per year. Under severe disease pressure, use the highest recommended rate and/or the shortest spray interval. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

NOTE: Do not use 26 GT Fungicide as a drench on impatiens, and pothos. Do not use 26 GT Fungicide on Spathiphyllum.

**HOW TO USE 26 GT FUNGICIDE AS A DIP**

PLANT SPECIES	DISEASE	QUARTS PRODUCT 100 GALLONS	DIP DURATION	DIRECTIONS
Rose	Botrytis Storage Rot (Botrytis sp.)	1.0	5 Minutes	Dip bare root roses prior to cold storage.
Azalea and Rhododendron	Cylindrocladium Blight and Wilt (Cylindrocladium scoparium)	1.0	5 Minutes	Dip cuttings prior to planting.
Gladiolus	Fusarium Corm Rot (Fusarium oxysporum)	2.0	5 Minutes	Dip corms prior to storage.

**TANK MIXTURES**  
Additional Disease Control: If ornamentals are threatened by additional diseases, 26 GT Fungicide is compatible with most commonly used fungicides. For control of diseases caused by Pythium and Phytophthora, 26 GT Fungicide can be tank-mixed with ALIETTE® brand fungicide. Consult the ALIETTE® label for a complete listing of diseases controlled and for application rates.

If a tank mix with other fungicides is used, follow label directions for the use of that product and apply at the recommended rate for control of the target disease organism.

**DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS**

- Do not use through sprinkler irrigation systems in California.
- Apply this product only through sprinkler irrigation systems including center pivot. Do not apply this product through any other type of irrigation system.

**SPRAY PREPARATION:** Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

**APPLICATION INSTRUCTIONS:** First prepare a suspension of 26 GT Fungicide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of 26 GT Fungicide, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of 26 GT Fungicide per 1 to 4 gallons of water are recommended) Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of 26 GT Fungicide into the irrigation water line so as to deliver the desired rate per acre. The suspension of 26 GT Fungicide should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with 26 GT Fungicide has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

**GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS**

- Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
- Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank

when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation must shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

**SPRAY DRIFT**

**SENSITIVE AREAS:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulation.

- The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

**INFORMATION ON DROPLET SIZE:** (This section is advisory in nature and does not supersede the mandatory label requirements)

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions below).

**CONTROLLING DROPLET SIZE:** (This section is advisory in nature and does not supersede the mandatory label requirements)

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**WIND:** (This section is advisory in nature and does not supersede the mandatory label requirements)

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY:** (This section is advisory in nature and does not supersede the mandatory label requirements)

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**TEMPERATURE INVERSIONS:** (This section is advisory in nature and does not supersede the mandatory label requirements)

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.


**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond Bayer Environmental Science's control, it is impossible for Bayer Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability or otherwise.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

© Bayer CropScience, 2006

CHIPCO, Compass and ProStar are registered trademarks of Bayer. 26 GT is a registered trademark of Bayer. Cleary's is a trademark of the Cleary Company. Daconil and Heritage are registered trademarks of Syngenta Corporation. Turfcide is a registered trademark of Chemtura Corporation.



Bayer Environmental Science  
A Business Group of Bayer CropScience LP  
Bayer Environmental Science  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709  
www.bayerprocentral.com

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING ▶

**Fungicide**  
A Fungicide for the Prevention and Control of Certain Diseases of Turfgrass and Ornamentals

**ACTIVE INGREDIENT:**  
Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide \* ..... 23.3%\*

**INERT INGREDIENTS:** ..... 76.7%

This product contains petroleum distillate.  
\*Equivalent to 2 pounds Iprodione per gallon.  
EPA Reg. No. 432-888 EPA Est. No. 264-MO-02

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For PRODUCT USE Information Call 1-800-334-9745.  
For MEDICAL and TRANSPORTATION Emergencies  
ONLY Call 24 Hours A Day 1-800-334-7577.

**NET CONTENTS**  
**2.5 gallons**  
4250930  
4588915 040618A

**BACKED  
by BAYER.**

**FIRST AID**

<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of medical emergency for additional information call toll free 1-800-334-7577.

**Note to Physician:** This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

**PRECAUTIONARY STATEMENTS  
CAUTION**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

**PERSONAL PROTECTIVE EQUIPMENT**

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment must wear long-sleeve shirt and long pants, chemical-resistant gloves such as barrier laminate, nitrile rubber (> 14 mils), neoprene rubber (> 14 mils), or viton (> 14 mils), chemical-resistant apron, and chemical-resistant footwear plus socks. Applicators using hand held equipment must wear coveralls over long-sleeve shirt and long pants, chemical-resistant gloves such as barrier laminate, nitrile rubber (> 14 mils), neoprene rubber (> 14 mils), or viton (> 14 mils), chemical-resistant footwear plus socks, chemical-resistant headgear for overhead exposures, and a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any R, P or HE filter. Applicators using mechanical ground equipment (groundboom, airblast, etc.) must wear long-sleeve shirt and long pants, and shoes plus socks. Applicators using truck-mounted equipment with a handgun at the end of a hose (i.e., for commercial turfgrass) and all other handlers not specified above must wear long-sleeve shirt and long pants, chemical-resistant gloves such as barrier laminate, nitrile rubber (> 14 mils), neoprene rubber (> 14 mils), or viton (> 14 mils), and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**ENGINEERING CONTROLS**

When handlers use closed systems, enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<p><b>User Safety Recommendations</b></p> <p>Users should:</p> <ul style="list-style-type: none"> <li>Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>
---

**ENVIRONMENTAL HAZARDS**

This chemical can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface water. This pesticide is toxic to invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**DIRECTIONS FOR USE**  
**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours for ornamental uses. The restricted entry interval for WPS uses is 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves such as barrier laminate, nitrile rubber (> 14 mils), neoprene rubber (> 14 mils), or viton (> 14 mils), and shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to turf uses (golf courses, landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter or allow others to enter the treated area until sprays have dried.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL**  
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL**  
Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

**General Use:** In order to assure maximum crop tolerance and disease control, follow recommendations on this label and all the precautions and limitations of the package label.

**GENERAL PRECAUTIONS AND RESTRICTIONS**

Use of this product at residential sites is prohibited. For use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application. For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present. Do not apply this product when the wind direction is toward aquatic areas.

**TURF**

26 GT® Fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. When used in conjunction with good turf management practices, 26 GT Fungicide is effective in controlling the following diseases:  
**Spring, Summer And Fall Diseases:** Dollar Spot, Brown Patch, Large Patch\*, Fusarium Blight and Necrotic Ring Spot\*, Leaf Spots such as Helminthosporium Leaf Spot caused by *Drechslera* spp. pathogens, Corticum Red Thread.  
**Winter Diseases:** Fusarium Patch (Pink Snow Mold) and Gray Snow Mold. Apply the recommended rates as indicated in the table in 0.5 to 10 gallons of water per 1000 square feet. Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

**TURF RECOMMENDATIONS FOR USE**

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

DISEASE	INTERVAL OF APPLICATIONS	FLUID OZ. 1000 FT.²
<b>Dollar Spot</b> ( <i>Lanzia</i> spp. and <i>Moellerodiscus</i> spp.)	<b>Greens and Tees:</b> Repeat at 14 to 21 day interval as long as required.	3 to 4
<b>Brown Patch</b> ( <i>Rhizoctonia solanii</i> ) <b>Leaf Spot</b> such as <i>Helminthosporium</i> Leaf Spot caused by ( <i>Drechslera</i> spp.)	<b>Fairways and Other Turf Areas:</b> Repeat at 14 to 28 day intervals as long as required.	<b>NOTE:</b> On Fairways, for Dollar Spot Control use 2 to 4 Fluid Ounces/1000 Ft.²
<b>Large Patch*</b> ( <i>Rhizoctonia solanii</i> )	Make first application in fall when conditions are favorable for disease development but no symptoms are visible. Make repeat applications in spring as needed on a 14-21 day interval.	4
<b>Fusarium Blight</b> ( <i>Fusarium</i> spp.) <b>Necrotic Ring Spot*</b> ( <i>Leptosphaeria korrae</i> )	Use only preventative foliar applications when conditions first become favorable for disease development. Additional applications should be made as necessary at 28 day intervals.	8
<b>Fusarium Patch</b> ( <i>Microdochium nivale</i> ) [Pacific Northwest Only – West of the Cascade Mountains]	Repeat at 14 to 21 day intervals as long as required.	4 to 8
<b>Gray Snow Mold</b> ( <i>Typhula</i> spp.) <b>Pink Snow Mold</b> ( <i>Fusarium nivale</i> )	One application before first permanent snow cover. If possible, another application during a mid-winter thaw.	4 to 8
<b>Corticium Red Thread</b> ( <i>Laetisaria luciformis</i> ) <b>Curvularia</b> ( <i>Curvularia</i> spp.)	Use as a preventative every 14 days as long as required. <i>Curvularia</i> control in bermudagrass only	4 4 to 8
<b>Anthracoese (suppression only)</b> ( <i>Colletotrichum</i> )	Will provide suppression during periods of anthracnose pressure; mix 26 GT with COMPASS or CHIPCO SIGNATURE or other anthracnose control fungicide	4 to 8

Do not exceed a total of 35 fluid oz. product/1000 ft² per year. Do not make more than 6 applications per year. Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

**TANK MIXTURES**

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred. Do not mix with any sticker, extender, or wetting agent. Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock or poultry. \*Not registered for use in California

**ADDITIONAL DISEASE CONTROL**

If turf is threatened by additional diseases, 26 GT Fungicide is compatible with most commonly used fungicides such as ProStar®, Compass®, and Heritage®. If a tank mixture is used, follow label directions for the use of that product. Do not exceed a total of 35 fluid oz. 26 GT Fungicide per 1000 ft² per year with a maximum of 6 applications. **Broad Spectrum Disease Control and Resistance Management:** A tank mixture of 26 GT Fungicide and Cleary's 3336 F provides effective, broad spectrum turf disease control and also serves as a useful tank mixture in the resistance management program required for other resistance sensitive fungicides.

Disease Pressure	26 GT +	Cleary's 3336 F
LOW	3 fl. oz./1000 sq. ft. +	1.0 fl oz./1000 sq. ft.
MEDIUM (more dollar spot and brown patch)	3 fl oz./1000 sq. ft. +	1.0 fl oz./1000 sq. ft.
HIGH	3 fl oz./1000 sq. ft. +	2.0 fl oz./1000 sq. ft.

**Summer Stress Complex/Summer Decline:** For management of Summer Stress Complex/Summer Decline, apply Chipco, Signature™ Fungicide at 4 to 8 ounces of product per 1000 square feet with 26 GT Fungicide at 2 to 4 ounces of product per 1000 square feet.

**Pythium Blight:** Pythium blight will be controlled by the tank mixing of CHIPCO® Signature™ Fungicide or ALIETTE® WDG brand Fungicide or Banol Fungicide with 26 GT Fungicide. If a tank mixture is used, follow label directions for the use of that product and apply at the rate recommended for control of the target disease organism.

**Gray Snow Mold:** For control of Gray Snow Mold (*Typhula* spp.) in areas where continuous snow cover occurs, refer to the following table for tank mixture recommendations.

Product	Disease	Rate
26 GT Fungicide	Gray Snow Mold	4-8 fl. oz.
Daconil 2787 Flowable	+	8 fl. oz.
Daconil 2787 WG	or	4.5 oz.
Daconil Ultrex 82.5 WG	or	4.9 oz.
Daconil Weather Stik 6F	or	5.5 fl. oz.
Turficide® 400	or	8 fl. oz.

Application must be made in autumn before snow cover occurs. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 square feet). Use the higher rate if the turf remains frozen prior to snow cover. For optimal control, reapply this treatment if a winter thaw and loss of snow cover occurs.

**ORNAMENTALS**

**NOT FOR RESIDENTIAL USE  
FIELD, LANDSCAPE AND GREENHOUSE ORNAMENTALS AND CONIFER NURSERIES**

CHIPCO® brand 26 GT Fungicide is a broad spectrum fungicide that may be applied safely to a wide range of ornamental flowering and foliage plants, either as a foliar spray, drench or dip. Please read specific instructions and use only as directed.

**RECOMMENDED FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL.**

CHIPCO® brand 26 GT Fungicide is recommended for use on a wide variety of flowering and foliage ornamentals as follows:

- |                                     |  |
|-------------------------------------|--|
| 1. Aerial Web Blight                | ( <i>Rhizoctonia</i> sp.)                                  |
| 2. Alternaria Leaf Blight           | ( <i>Alternaria euphorbiae</i> )                           |
| 3. Alternaria Leaf Spot             | ( <i>Alternaria panax</i> , <i>Alternaria tenuissima</i> ) |
| 4. Botrytis Blight                  | ( <i>Botrytis</i> sp.)                                     |
| 5. Fusarium Leaf Spot               | ( <i>Fusarium moniliforme</i> )                            |
| 6. Helminthosporium Leaf Spot       | ( <i>Helminthosporium</i> sp.)                             |
| 7. Rhizoctonia stem and root rot    | ( <i>Rhizoctonia</i> sp.)                                  |
| 8. Ink Spot                         | ( <i>Drechslera iridis</i> )                               |
| 9. Tulip Fire                       | ( <i>Botrytis tulipae</i> )                                |
| 10. Alternaria Leaf Blight          | ( <i>Alternaria zinniae</i> )                              |
| 11. Ray Blight                      | ( <i>Ascochyta chrysanthami</i> )                          |
| 12. Fusarium Corm rot               | ( <i>Fusarium oxysporum</i> )                              |
| 13. Daffodil Leaf Scorch            | ( <i>Stagnospora curtisii</i> )                            |
| 14. Blossom Blight                  | ( <i>Monilinia fructicola</i> )                            |
| 15. Botrytis Storage Rot            | ( <i>Botrytis</i> sp.)                                     |
| 16. Cylindrocladium Blight and Wilt | ( <i>Cylindrocladium scoparium</i> )                       |

**PLANT TOLERANCE:** Plant tolerances to 26 GT Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to 26 GT Fungicide. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

**ORNAMENTALS**

- |                              |                                 |                        |
|------------------------------|---------------------------------|------------------------|
| Ageratum (1 to 7)            | Chrysanthemum (1 to 7, 11)      | Dianthus (1 to 7)      |
| Ajuga (1 to 7)               | Cineraria (1 to 7)              | Dieffenbachia (1 to 7) |
| Almond (ornamental) (1 to 7) | Cistena Plum (1 to 7, 14)       | Dizygotheca (1 to 7)   |
| Alyssum (1 to 7)             | Coleus (1 to 7)                 | Dogwood (1 to 7)       |
| Andromeda (1 to 7)           | Columbine (1 to 7)              | Dracena (1 to 7)       |
| Aphelandra (1 to 7)          | Coral Bells (Heuchera) (1 to 7) | English Ivy (1 to 7)   |
| Artemisia (1 to 7)           | Crape Myrtle (1 to 7)           | Episcia (1 to 7)       |
| Aster (1 to 7)               | Crassula (1 to 7)               | Euonymous (1 to 7)     |
| Azalea (1 to 7, 16)          | Croton (1 to 7)                 | Ficus (1 to 7)         |
| Boxwood (1 to 7)             | Cyclamen (1 to 7)               | Forsythia (1 to 7)     |
| Cactus (1 to 7)              | Daffodils (1 to 7, 13)          | Gazania (1 to 7)       |
| Calendula (1 to 7)           | Dahlia (1 to 7)                 | Geranium (1 to 7)      |
| Carnation (1 to 7)           | Delphinium (1 to 7)             | Gladiolus (1 to 7, 12) |
| Cherry (ornamental) (1 to 7) | Deutzia (1 to 7)                | Gloxinia (1 to 7)      |

# 26 GT™ Fungicide

A Fungicide for the Prevention and Control of Certain Diseases of Turfgrass and Ornamentals

## ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide\* . . . . . 23.3%\*

**INERT INGREDIENTS:** . . . . . 76.7%

This product contains petroleum distillate. \*Equivalent to 2 pounds Iprodione per gallon.

EPA Reg. No. 432-888

EPA Est. No. 264-MO-02

## KEEP OUT OF REACH OF CHILDREN CAUTION

For **PRODUCT USE** Information Call 1-800-334-9745.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577.

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface water.

This pesticide is toxic to invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

For Additional Precautionary Statements and Complete Directions For Use, See Attached Booklet.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of medical emergency for additional information call toll free 1-800-334-7577.

**Note to Physician:** This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

### PRECAUTIONARY STATEMENTS

#### CAUTION

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

#### ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential



Bayer Environmental Science

A Business Group of  
Bayer CropScience LP

Bayer Environmental Science  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709  
www.bayerprocentral.com

**NET CONTENTS: 2.5 gallons**

**4250930**

4588915 040618A

PostScriptPicture

(Code 128, 0100785740862377 . eps)