



# PERMANONE® 10% E.C.

\* RECOMMENDED FOR AGRICULTURAL/COMMERCIAL USE

\* BROAD SPECTRUM MULTI-USE INSECTICIDE

\* QUICK KNOCKDOWN

\* FOR USE ON LIVESTOCK: CATTLE (BEEF AND DAIRY), GOATS, SHEEP, SWINE, HORSES AND POULTRY

ACTIVE INGREDIENT:

\*Permethrin: . . . . . 10.0%

†OTHER INGREDIENTS: . . . . . 90.0%

100.0%

\*Cis/trans ratio: Max. 65% (+/-) trans and min. 35% (+/-) cis.

†Contains Petroleum Distillates

EPA Reg. No. 432-1132

EPA Est. No. 432-TX-1

## KEEP OUT OF REACH OF CHILDREN WARNING

FIRST AID	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER: 1-800-331-2867</b>	
[Synthetic Pyrethroid]	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.	
<b>Note To Physician:</b> Contains petroleum distillate – vomiting may cause aspiration pneumonia. Do not administer or instill milk, cream or other substances containing vegetable or animal fats which enhance absorption of lipophilic substances.	

See Side Panel For Additional Precautionary Statements.

**NET CONTENTS: 2.5 gallons**

**READ ENTIRE LABEL BEFORE EACH USE**

**PRECAUTIONARY STATEMENTS**  
**Hazards To Humans & Domestic Animals**

**WARNING**

Causes skin irritation. Do not get on skin or on clothing. Wear coveralls worn over short-sleeved shirt and short pants, socks, chemical resistant foot wear and gloves when handling or mixing concentrate. Harmful if inhaled or absorbed through skin. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**Environmental Hazards**

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when wind speeds exceed 10 mph. Do not contaminate water when disposing of equipment washwaters. Shrimp and crab may be killed if application rates recommended on this label are exceeded. Use with caution where these are important resources. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply this product or allow drift when bees are actively visiting the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**Physical & Chemical Hazards**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To prepare dilutions, the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water or oil and blend thoroughly. Do not hold dilutions for more than 24 hours.

Do not allow spray treatment to contaminate pasture land, cropland, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture

APPLICATION SITES	PEST CONTROLLED	USE DIRECTIONS	DILUENT	DILUTION RATE	APPLICATION RATE	DOSE RATE
LIVESTOCK SPRAY: Dairy & Beef Cattle, Sheep, Goats, Swine, Horses and Poultry. (For swine, do not ship animals for slaughter within 5 days of last treatment.)	Horn Flies, House Flies, Mosquitoes, and Gnats	Apply to wet the hair thoroughly, with particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.	Water	1 part concentrate in 100 parts water - (1.3 fl. ounces per gallon)		
	Stable Flies, Horse Flies and Deer	Apply a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatment as needed.	Water	1 part concentrate in 100 parts water - (1.3 fl. ounces per gallon)		
	Face Flies	Apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before release to pasture. Apply sufficiently to wet the face but not more than 1.5 ounces per animal. Repeat daily as needed.	Water	1 part concentrate in 100 parts water (1.3 fl. ounces per gallon)		
	Sheep "ticks" or keds	Thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.	Water	1 part concentrate in 100 parts (1.3 fl. ounces per gallon)		
	Fleas on Livestock (To kill and to protect against reinfestation)	Wet the animal by dipping or spraying.	Water	1 part concentrate in 20 parts water (6.5 fl.ounces per gallon)		
AS A SPACE SPRAY Animal Houses, Corrals, Drive-in restaurants, Drive-in theaters, Feedlots, Gardens, Golf courses, Parks, Playgrounds, Recreational Areas, Urban Areas, Zoos, etc.	Mosquitoes, Gnats, biting and non-biting Midges, Black- flies, and other biting Flies.	Apply using vehicle mounted ULV equipment to create an insecticidal swath. For best results apply when there is a light breeze (ie. about 5 mph) and apply in direction of breeze to obtain maximum swath width and better distribution. Do not apply if winds exceed 10 mph.  The target dose rate to be applied is up to 0.007 lb. a.i. per acre. This can be achieved under different conditions by altering the flow of insecticide from the equipment and/or the speed of the vehicle. Apply 3.5 to 7.0 fluid ounces per minute at vehicle speeds of 5-10 miles per hour using	Oil or Wateroil	1 part concentrate with 1 part suitable solvent, or at proportional oil dilutions calculated on the basis of vehicle speed and swath width to achieve the target dose.	Dependent on swath width, flowrate and speed of vehicle.	Up to 0.007 lbs of active ingredient per acre.

		<p>swath widths of 150-300 feet as appropriate to local conditions.</p> <p>At higher vehicle speeds, flow rates should be increased proportionately to achieve the target dose rate.</p> <p>Not to be used within 100 feet (30 meters) of lakes and streams.</p> <p>In treatment of corrals, feed lots and swine lots, cover any exposed drinking water, drinking fountains and animal feed before application.</p> <p>Allow 24 hours before retreating. Do not apply more than once in a 24 hour period.</p>				
<p><b>BARRIER SPRAY</b> Perimeter of residential yards, public facilities, paths and recreational areas.</p>	<p>As an aid in reducing mosquito annoyance. To kill and control biting and non-biting midges, deer flies.</p>	<p>Apply when walking at a speed of approximately 2 mph or a swath of 50 feet treating approximately 0.2 acre/minute.</p>	<p>Oil or Water</p>	<p>1 part concentrate to 1 part of a suitable diluent, or at proportional dilutions calculated on the basis of applicator speed and swath width to achieve the target dose.</p>	<p>Apply 17.5 fluid ounces of diluted material per acre.</p>	<p>Approx. 0.1 lb per acre.</p>

### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage And Spill Procedures: Store upright at room temperature.

Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

CONTAINERS LARGER THAN ONE GALLON: Metal Containers – Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers - Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Glass Containers - Triple rinse or equivalent, then dispose of in a sanitary landfill or by other approved state and local procedures.

### **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer Environmental Science. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©BayerCropScience LP, 2006

PERMANONE is a registered trademark of Bayer.



**Bayer Environmental Science**

Bayer Environmental Science  
A Business Group of Bayer CropScience LP  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709

102604